START THE TALK. SEE THE DIFFERENCE.

Dry eye disease is a common condition that can significantly impact your daily life.

vevye® (cyclosporine ophthalmic solution) 0.1%

Complete the questions below and discuss your results with your doctor to determine if you may have dry eye disease, and if VEVYE® *(cyclosporine ophthalmic solution) 0.1%* is the right choice for you.

SYMPTOMS

What symptoms are you experiencing? Note the severity and frequency.

CURRENT MANAGEMENT

List any treatments or remedies you are currently using (e.g., artificial tears, warm compresses). Note any relief or side effects from these treatments.

IMPACT ON DAILY LIFE

Describe how dry eye affects your daily activities (e.g., reading, driving, looking at screens, outdoor activities).

INDICATION AND USAGE

- VEVYE® (cyclosporine ophthalmic solution) 0.1% is indicated to treat the signs and symptoms of dry eye disease.

IMPORTANT SAFETY INFORMATION

- To avoid the potential for eye injury and/or contamination, do not touch the bottle tip to the eye or other surfaces.
- VEVYE should not be administered while wearing contact lenses. Remove contacts prior to using VEVYE. After applying VEVYE to the eyes wait 15 minutes before reinserting contact lenses.
- The most common side effects include eye irritation (8%) and temporary blurred vision (3%).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For additional information about VEVYE, go to vevye.com.

VEVYE, and the VEVYE logo are registered trademarks of Novaliq GmbH.
Harrow and the Harrow logo are trademarks of Harrow IP, LLC.

© 2024 Harrow. All Rights Reserved. VYE-00140 10/24

HARROW

Is VEVYE® *(cyclosporine ophthalmic solution) 0.1%* right for you?

It's important to consult your doctor to determine if VEVYE® is the best option for you.

vevye®
(cyclosporine ophthalmic solution) 0.1%

Here are some questions to consider asking about VEVYE:

- **How does VEVYE differ from what I'm currently using?**
- **How often do I use VEVYE?**
- **How long will a bottle of VEVYE last?**
- **What are the most common side effects?**
- **Does VEVYE have a Savings Program?**
- **How long will I need to continue using VEVYE?**

NOTES

INDICATION AND USAGE

- VEVYE® (cyclosporine ophthalmic solution) 0.1% is indicated to treat the signs and symptoms of dry eye disease.

IMPORTANT SAFETY INFORMATION

- To avoid the potential for eye injury and/or contamination, do not touch the bottle tip to the eye or other surfaces.
- VEVYE should not be administered while wearing contact lenses. Remove contacts prior to using VEVYE. After applying VEVYE to the eyes wait 15 minutes before reinserting contact lenses.
- The most common side effects include eye irritation (8%) and temporary blurred vision (3%).

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088.

For additional information about VEVYE, go to vevye.com.

VEVYE, and the VEVYE logo are registered trademarks of Novaliq GmbH.
Harrow and the Harrow logo are trademarks of Harrow IP, LLC.

© 2024 Harrow. All Rights Reserved. VYE-00140 10/24

HARROW